

for Animal use only

Xinanning

【Scientific name/ Generic Name】 Newcastle Disease Vaccine, Live (Strain La Sota).

【Trade Name】 Xinanning.

【Active Component】 Live attenuated Newcastle Disease Virus, La Sota Strain $\geq 10^{6.0}$ EID₅₀/dose.

【Physical properties】 The product is a kind of spongiform loose mass. It can be separated easily from the vial wall, and quickly dissolve in diluents.

【Pharmaceutical form】 live attenuated, freeze dried vaccine.

【Indication / Effect】 The product is used for protection from Newcastle Disease in chickens.

【Dosage and Administration】

Product can be applied via Nasal, Ocular, Oral or Respiratory Routes

The product can be used for chickens more than 7 days old; booster vaccinations can be applied according to your veterinarian advice.

Nasal and ocular routes: Each 1000 dose vial should be diluted in 30 ml sterilized physiological saline or injection water or distilled water, then apply 1 drop each (0.03ml)/bird.

Drinking water route, the vaccine is diluted in drinking water, water volume is determined according to Bird age for ex 5~10 days old, 10 ml each; 20~30 days old, 10~20ml each; adult chicken, 20~30ml each. Generally, dissolve the vaccine in a volume of drinking water that can be ingested within 1 hour.

Respiratory route (spraying): fill the apparatus with fresh tap water and spray the surface which birds occupy, in a manner so that the heads of all the birds are covered with droplets of water. Check the amount of water used, and this will be used to mix with the necessary doses, depending on the number of birds to be vaccinated. The following volumes are guidelines:

| No. of doses | chicks less than 30 days (large droplets) | birds older than 30 days (fine droplets) |
|--------------|---|--|
| 1,000 | 200-250 ml | 500-1,000 ml |

Please refer to your veterinarian to establish the most suitable vaccination program according to the health conditions of each farm and area.

北京信得威特科技有限公司 Beijing Sinder-vet Technology Co., Ltd.

A subsidiary of Shandong Sinder Technology Co., Ltd.

Add.: No.118, Shunyu Road, Beijing Tianzhu Airport Economic Development Zone, Shunyi District, Beijing, China.

Tel: [+86 532 58820810](tel:+8653258820810)

Email: alice@sindergroup.cn

Website: www.sinder.cn

【Precautions】

1. It is forbidden to use spraying application if there is mycoplasma infection in chicken flocks.
2. Store the product in cool, dark place and should be used up in 4 hours after dilution.
3. For oral vaccination, drinking water should be clean and contain no disinfectants such as chlorine. Don't use metal container for dilution.
4. After use, empty vaccine vials, unused vaccine and disposable appliances should be safely disposed of.
- 5- Keep away from Heat and direct sunlight.

【Withdrawal period】 : Zero days.

【Presentations】 1000 doses/vial. 2000 doses/vial. 5000 doses/Vial.

【Package】 10 vials/Foam box × 30 Foam Boxes/Carton.

【Storage】 Store at 2-8°C, Shelf life is 20 months from production date.

【Drug Approval No.】 Shouyaoshengzi 010372007.

【Manufacturer】 Beijing Sinder-vet Technology Co., Ltd., China (A Subsidiary of Shandong Sinder Technology Co. LTD), China.

【Exporter】 Shandong Sinder Technology Co., Ltd., China.

Sole Agent in Egypt:

Egyptian Marketing and Distribution Center

28 Murad Street, Giza, Egypt.

Phone 02/ 35725272 Fax: 02/53721668