

Leaflet**For vet. use**

## **XINNUONING**

*Combined Newcastle Disease & Infectious Bronchitis Vaccine, Live*

*(Strain HB1 + Strain H120) for Chickens*

### **【Scientific Name / Veterinary Drug Name】**

Combined Newcastle Disease & Infectious Bronchitis Vaccine, Live (strain HB1 + strain H120).

### **【Trade Name】XINNUONING.**

### **【Main Component and Content】**

Each one dose contains:  $\geq 10^{3.5}$  EID<sub>50</sub> Infectious Bronchitis virus, strain H120 &  
 $\geq 10^6$  EID<sub>50</sub> Newcastle Disease virus, strain Hitchner B1 (HB1).

### **【Pharmaceutical form】** Bivalent, live attenuated, freeze dried vaccine.

**【Physical properties】** a faint yellow to slight reddish spongiform loose mass. It can be easily separated from the vial wall, and quickly dissolve in diluents.

**【Usage】** For active immunization against Newcastle Disease and Infectious Bronchitis in chickens.

### **【Dosage and Administration】**

**Routes of Administration:** Nasal dropping or drinking water vaccination.

The product is used for chickens more than 7 days old; booster dosage can be applied according to your veterinarian advice.

**Nasal dropping:** Each 1000 dose vial will be diluted in 30 ml sterilized physiological saline or injection water or distilled water, then apply 1 drop (0.03ml)/bird.

**Drinking water vaccination,** the vaccine is diluted in drinking water, water volume is determined according to Bird age for ex 5~10 days old, 10 ml each; 20~30 days old, 10~20ml each; adult chicken, 20~30ml each. Generally, dissolve the vaccine in a volume of drinking water that can be ingested within 1 hour.

Please refer to your veterinarian to establish the most suitable vaccination program according to the health conditions of each farm and area.

### **【Precautions】**

1. After dilution the vaccine should be kept in dark cool place.
2. Metal containers shouldn't be used for product dilution.
- 3- Don't use water with chlorine or disinfectants.

北京信得威特科技有限公司 **Beijing Sinder-vet Technology Co., Ltd.**

**A subsidiary of Shandong Sinder Technology Co., Ltd.**

**Add.: No.118, Shunyu Road, Beijing Tianzhu Airport Economic Development Zone, Shunyi District, Beijing, China.**

**Tel: +86 532 58820810**

**Email: [alice@sindergroup.cn](mailto:alice@sindergroup.cn)**

**Website: [www.sinder.cn](http://www.sinder.cn)**

- 4- Don't expose to heat or direct sunlight.
- 5- In case of oral vaccination, don't allow birds to drink water at 1~4 hours before vaccination.
6. Vaccine vials, implement, unspent vaccine must be disposed hygienically.
- 7- Ensure adequate hygiene and management during post-vaccination period.

**【Presentations】** 1000 doses/vial, 2000 doses/vial, 5000 doses/vial.

**【Package】** 10 vials/ Box × 30 Boxes/Carton.

**【Withdrawal period】** : Zero days.

**【Storage】** Store at 2-8°C.

**【Shelf life】** 18 months from production date.

**【Drug Approval No.】** Shou Yao Sheng Zi 010372021

**【Manufacturer】** Beijing Sinder-vet Technology Co., Ltd., China, " A subsidiary of Shandong Sinder Technology Co., Ltd", China.

**【Exporter】** Shandong Sinder Technology Co., Ltd ,China.

**Sole Agent in Egypt:**

**Egyptian Marketing and Distribution Center**

28 Murad Street, Giza, Egypt.

Phone 02/ 35725272 Fax: 02/53721668

北京信得威特科技有限公司 **Beijing Sinder-vet Technology Co., Ltd.**

**A subsidiary of Shandong Sinder Technology Co., Ltd.**

**Add.: No.118, Shunyu Road, Beijing Tianzhu Airport Economic Development Zone, Shunyi District, Beijing, China.**

**Tel: [+86 532 58820810](tel:+8653258820810)**

**Email: [alice@sindergroup.cn](mailto:alice@sindergroup.cn)**

**Website: [www.sinder.cn](http://www.sinder.cn)**