
Sinder ND-K

Inactivated Newcastle Disease Vaccine





Another Source of Reliable Poultry Vaccines & Biologicals

Remove Watermark Now

- Totaly Safe For Young Chickens.
- Provide Early and long lasting Protection.
- Low Viscosity adjuvant is well Tolerated at Injection Site.
- Exceptional Serological Response.

Indication:

For vaccination of Chickens against Newcastle disease in areas where highly pathogenic ND is endemic.

Composition:

ND inactivated Virus, Pre-inactivation Titer ≥10⁸ EID 50/ 0.5 ml Dose.

Pharmaceutical form: Water in oil Emulsion.

Posology and Adminstartion:

Broilers: 0.25 ml dose S/C during first week of life ,Breeders and Layers: initial dosage 0.25 ml S/C during first week of life ,boosted by 0.5 ml dosage 10 weeks from initial dosage Followed by 0.5 ml

dosage S/C or I.M about two weeks before laying.

Storage: Store at 8-2 C. Dont Freeze.

Withdrawal Time: 28 Days.

Adverse Effects: Generally there aren't any adverse effects if product is used properly.

Packing: 250 ml - 500 ml HDPE Bottles.

* Consult your Veterinarian about the proper vaccination program for your area.



EMDC



Sole Agent:

Egyptian Marketing and Distribution Center 28 Morad Street, Giza.

Phone: 35725272-02 35728355-02.

SINDER

Beijing Sinder-Vet Technology Co.,Ltd. Beijing- PRC, www.sinder.cn (A Sumitomo Joint Venture Company)